

Journal of Asia-Pacific Biodiversity Vol. 6, No. 2 249-253, 2013
<http://dx.doi.org/10.7229/jkn.2013.6.2.249>

Journal of
Asia-Pacific
Biodiversity

The Fauna of Ground Beetles (Coleoptera: Carabidae) in Paddy Fields, Four Province, Korea

Jinyoung Park¹, Seong Joon Park² and Jong Kyun Park^{1*}

¹Department of Applied Biology, College of Ecology & Environmental Sciences, Kyungpook National University, Sangju 742-711, Republic of Korea

²National Institute of Environmental Research, Incheon-si, 404-708, Republic of Korea

Abstract: A total of 29 species belonging to 17 genera of 10 subfamilies were identified from 277 collected ground beetles in paddy fields. [Species richness was high in Harpalinae (9 species, 31.0%), Pterostichinae (8 species, 27.6%), Callistinae (4 species, 13.8%), Nebriinae (2 species, 6.9%), and others subfamilies (1 species, 3.4%). The subfamily Harpalinae had the most number of individuals (24.5%), followed by 63 Callistinae (22.7%), 59 Zabrinae (21.3%), and 48 Brachininae (17.3%), 19 Pterostichinae (6.9%), 8 Licininae (2.9%), Nebriinae (2.5%), 3 Patrobinae (1.1%), and 1 other subfamily (0.4%), respectively. The dominant species was *Amara congrua* (59 individuals, 21.3%) and the subdominant species was *Chlaenius pallipes* (49 individuals, 17.7%). Between Site 4 (Chuncheon-city, Gangwon-do) and Site 5 (Gangneung-city, Chuncheon-city, Gangwon-do) and between Site 1 (Yuseong-gu, Daejeon metropolitan-city) and Site 2 (Danyang-gun, Chungcheongbuk-do) had the highest similarity in the result of cluster analysis using the surveyed ground beetles.

Keywords: Carabidae, Monitoring, Inventory, Diversity, Paddy Fields, Field

Introduction

A paddy field is a flooded parcel of arable land used for growing rice and other semi-aquatic crops. Paddy fields account for approximately 15% of the global arable land area (Huke and Huke, 1997, Maclean *et al.* 2002). In South Korea, paddy fields are the most common type of agricultural land, covering approximately 60% of all agricultural land (Do *et al.*, 2011).

Paddy fields provide forage, refuge, and a place for overwintering or estivation for many species, including soil macrofauna and microfauna, insects, and birds (Kato, 2001, Thomas *et al.*, 2004, Katoh *et al.*, 2009, Paik *et al.*, 2009). Among these, ground beetles except Harpalinae and Zabrinae, are predaceous and a natural enemy of small-sized invertebrates including earthworms, aphids, moths and snails, that play a very important role in the ecosystem (Lövei and Sunderland, 1996; Holland, 2002) and this considers ground beetles an important natural enemy group in mountainous areas and agricultural environment (Kromp, 1999; Holland, 2002). More specifically, in wet habitats such as paddy fields, abandoned paddy fields, wet grasslands, riversides, and lowlands with different vegetation, lower

soil pH, and higher soil moisture than surrounding areas, ground beetles can be characterized by species composition, food preference, and habitat selection (Hengeveld, 1987, Luff *et al.*, 1989, Eyre *et al.*, 1990, Do and Moon, 2002, Do *et al.*, 2007). Some of these studies have examined the effectiveness of restoring abandoned paddy fields with respect to improved agricultural landscape biodiversity (Comin *et al.*, 2001, Yamada *et al.*, 2007, Uematsu *et al.*, 2010).

The study was performed to make specific inventories of ground beetles in paddy fields and to provide fundamental information and diversity on community structure of ground beetles.

Materials and Methods

Survey Sites and Collecting Method

Ground beetles mainly live on the surface of the paddy fields and pitfall traps are installed considering these features. There are 10 traps at 10 m interval for each and the top of the trap is placed at the same height of the surface of the paddy field. Transparent plastic bottles with 10.0 cm height, 7 cm diameter and 200 mL volume were used as traps and had plastic filters with 6 holes at 2 cm diameter to prevent from mid- and large-sized animals like rodents. Traps were filled with preservatives (50 mL, environmentally friendly antifreeze, Super-A Green, SK chemicals, Korea).

*To whom correspondence should be addressed.
Tel: +82-54-530-1215; Fax: +82-54-530-1218
E-mail: entopark@knu.ac.kr




Fig. 1. The surveyed sites of ground beetles in Korea.

Pitfall traps were collected approximately every four weeks between May to July-2011.

We surveyed at the following five sites according to the administrative district in Korea (Fig. 1);

Site 1. Seongbuk-dong, Yuseong-gu, Daejeon metropolitan-city

Site 2. Daedae-ri, Gagok-myeon, Danyang-gun, Chungcheongbuk-do

Site 3. Sinsong-ri, Seocheon-eup, Seocheon-gun, Chungcheongnam-do

Site 4. Yeochan-ri, Gujeong-myeon, Gangneung-city, Gangwon-do

Site 5. Bangdong-ri, Seo-myeon, Chuncheon-city, Gangwon-do

Identification

Collected ground beetles were brought to the laboratory and dried, mounted, and identified to the species level under a dissecting microscope (Olympus, SZ40, $\times 20$). Identification was performed according to Habu (1967, 1973, 1978), Kwon and Lee (1984), and Park and Paik (2001), Löbl and Smetana (2003), and Park (2004). The specimens examined were deposited in the J.Y. Park Collection (JYPC).

Community structure analysis

Pielou's species diversity index (H' , Pielou, 1969), McNaughton's dominance index (DI, McNaughton, 1967), Margalef's species richness index (RI, Margalef, 1958), Pielou's species evenness index (EI, Pielou, 1975) and Jaccard's similarity index (Jaccard, 1908) were calculated and the formulas are as follows:

$$H' \text{ (Species diversity index)} = - \sum \left[\frac{n_i}{N} \cdot \log_2 \frac{n_i}{N} \right]$$

n_i means number of individuals at i -th species and N means total number of individuals (Pielou, 1969).

$$DI \text{ (Dominance index)} = \frac{n_1 + n_2}{N}$$

n_1 means number of dominant species individuals, n_2 means number of subdominant species individuals, N means total number of individuals (McNaughton, 1967).

$$RI \text{ (Species richness index)} = \frac{S - 1}{\ln(N)}$$

S means total number of species and N means total number of individuals (Margalef, 1958).

$$EI \text{ (Evenness index)} = \frac{H'}{\log_2 S}$$

H' means species diversity index and S means total number of species (Pielou, 1975).

Results



A total of 29 species belonging to ten subfamilies were identified from 277 collected ground beetles in paddy fields (Table 1). Nine species of Harpalinae recorded the highest number of subfamily species, followed by 8 Pterostichinae, 4 Callistinae, 2 Nebrinae and 1 other subfamily, respectively. The subfamily Harpalinae had the most number of individuals (24.5%), followed by 63 Callistinae (22.7%), 59 Zabrinae (21.3%), 48 Brachininae (17.3%), 19 Pterostichinae (6.9%), 8 Licininae (2.9%), Nebrinae (2.5%), 3 Patrobinae (1.1%) and 1 other subfamily (0.4%), respectively (Fig. 2, 3).

At the genus level, 5 species of *Harpalus*, 4 species of *Chlaenius*, 2 species of *Anisodactylus*, *Nebria*, *Colpodes* and *Pterostichus* genus were collected. Also, the other 11 genera were for 1 species, respectively (Fig. 4). At the genus level, 63 individuals of *Chlaenius* and 59 individuals of *Amara* genus were collected, followed by *Pheropsophus* and *Harpalus* with 48 and 43, respectively (Fig. 5). The number of ground beetles species in each surveyed site were from 6 (Site 1, 3 Yuseong-gu, Daejeon metropolitan-city, Seocheon-gun, Chungcheongnam-do) to 12 (Site 2 Danyang-gun, Chungcheongbuk-do, Fig. 6). The dominant species was *Amara congrua* (59 individuals, 21.3%) and the subdominant species was *Chlaenius pallipes* (49 individuals, 17.7%). The Dominance index (DI) for each sites were 0.45 to 0.88, and the average dominance index was in the order of St. 3 > St. 5 > St. 1 > St. 4 > St. 2, respectively.

The species diversity index (H') for each sites were 1.26

Table 1. List of ground beetles collected in Paddy fields. Wing form: B, brachypterous; M, macropterous

Subfamily	Species	Wing form	St.1	St.2	St.3	St.4	St.5
Harpalinae	<i>Harpalus chalcatus</i>	M		14			4
	<i>Harpalus tridens</i>	M		3			
	<i>Harpalus jureceki</i>	M					3
	<i>Harpalus bungii</i>	B	5	1			
	<i>Harpalus crates</i>	M		13			
	<i>Oxycentrus argutoroides</i>	M				3	
	<i>Anoplogenus cyaneus</i>	M			7		
	<i>Anisodactylus signatus</i>	M		2	2		
	<i>Anisodactylus punctatipennis</i>	M	4	7			
Patrobinae	<i>Patrobis flavipes</i>	M			3		
Callistinae	<i>Chlaenius ocreatus</i>	M				1	
	<i>Chlaenius naeviger</i>	M				4	8
	<i>Chlaenius circumductus</i>	M					1
	<i>Chlaenius pallipes</i>	M	2	2		2	43
Lebiinae	<i>Calleida onoha</i>	M			1		
Pterostichinae	<i>Poecilus nitidicollis</i>			3			
	<i>Pterostichus subovatus</i>	M			1		
	<i>Pterostichus microcephalus</i>						4
	<i>Dolichus halensis halensis</i>	B				1	
	<i>Synuchus congruus</i>	M				2	
	<i>Colpodes xestus</i>	B		3			
	<i>Colpodes japonicus</i>						4
	<i>Curtonotus giganteus</i>		1				
Zabrinae	<i>Amara congrua</i>	M	8	1	44		6
Nebriinae	<i>Nebria chinensis</i>	M	2				
Nebriinae	<i>Nebria coreica</i>	M					5
Carabinae	<i>Calosoma chinense</i>	B		1			
Brachininae	<i>Pheropsophus jessoensis</i>	B		15		6	27
Licininae	<i>Lachnocrepis japonica</i>	M				8	


**Fig. 2.** Species ratio of each taxon.**Fig. 3.** Individual ratio of each taxon.

to 2.98, and the average species diversity index was in the order of St. 2 > St. 4 > St. 5 > St. 1 > St. 3, respectively.

The species richness index (R') for each sites were 1.23 to 2.64, and the average species diversity index was in the order of St. 2 > St. 4 > St. 5 > St. 1 > St. 3, respectively.

The species evenness index (E') for each sites were 0.49 to 0.89, and the average species diversity index was in the order of St. 4 > St. 1 > St. 2 > St. 5 > St. 3, respectively (Table 2).

Between Site 4 (Chuncheon-city, Gangwon-do) and Site

**Fig. 4.** The number of species of each genus.




Fig. 5. The number of individuals of each genus.




Fig. 6. The number of species and individuals in each surveyed site.

Table 2. Various community analysis of each surveyed site

Site	Num. Sp./Indivi.	DI	H'	R'	E'
St. 1	6/22	0.59	2.29	1.62	0.89
St. 2	12/65	0.45	2.98	2.64	0.83
St. 3	6/58	0.89	1.26	1.23	0.49
St. 4	8/27	0.52	2.67	2.12	0.89
St. 5	10/105	0.67	2.51	1.93	0.76
Mean	29/277	0.62	2.34	1.91	0.77

Sp: Number of species; Indivi: Number of individuals; DI: Dominance index; H': Species diversity; R': Species richness index; E': Evenness index




Fig. 7. The cluster analysis of collected ground beetles in each surveyed site.

5 (Gangneung-city, Chuncheon-city, Gangwon-do) and between Site 1 (Yuseong-gu, Daejeon metropolitan-city) and Site 2 (Danyang-gun, Chungcheongbuk-do) had the similarity in the result of cluster analysis using the surveyed ground beetles (Fig. 7).

Acknowledgments

This study was supported by and carried out under the financial support of National Institute of Environmental Research (NIER).

References

Comimn FA, Romero JA, Hernández O and Menéndez M. 2001. Restoration of wetlands from abandoned rice fields for nutrient removal, and biological community and landscape diversity. *Restor. Ecol.* 9: 201-208.

Do YH, Moon TY. 2002. Succession of insect communities by desiccation of bog Hwaemneup at Mt. Wonhyosan, Yangsan. *J. Korean Wetl. Soc.* 4: 13-22. (in Korean with English summary)

Do YH, Moon TY, and Joo GJ. 2007. Application of the carabid beetles as ecological indicator species for wetland characterization and monitoring in Busan and Gyeongsangnam-do. *Korean J. Environ. Ecol.* 21: 22-29. (in Korean with English summary)

Do YH, Jeong KS M, Lineman JY, Kim HA, and Joo GJ. 2011. Community changes in carabid beetles (Coleoptera: Carabidae) through ecological succession in abandoned paddy fields. *J. Ecol. Field Biol.* 34(3): 269-278.

Eyre MD, Luff ML, and Rushton SP. 1990. The ground beetle (Coleoptera, Carabidae) fauna of intensively managed agricultural grasslands in northern England and southern Scotland. *Pedobiologia.* 34: 11-18.

Habu A. 1967. *Fauna Japonica, Carabidae truncatipennes group* (Insecta: Coleoptera). Biogeographical Society of Japan. Tokyo. Japan. 338 pp. plates.

Habu A. 1973. *Fauna Japonica, Carabidae: Harpalini* (Insecta: Coleoptera). Keigaku Publishing Co. Ltd. Tokyo. Japan. 430 pp. 21 plates.

Habu A. 1978. *Fauna Japonica, Carabidae: Platynini* (Insecta: Coleoptera). Keigaku Publishing Co. Ltd. Tokyo. Japan. 447pp. 36 plates.

Hengeveld R. 1987. Scales of variation: their distinction and ecological importance. *Ann. Zool. Fenn.* 24: 195-202.

Holland JM. 2002. *The agroecology of carabid beetles.* InterceptLtd. Andover, Hampshire, UK. 356pp.

Huke RE, Huke EH. 1997. *Rice Area by Type of Culture: South, Southeast, and East Asia maps.* International Rice Research Institute, Manila.

Jaccard P. 1908. Nouvelles recherches sur la distribution florale. *Bull. Soc. Vaud. Sci. Nat.* 44: 223-207.

Kato M. 2001. 'Satoyama' and biodiversity conservation: 'Satoyama' as important insect habitats. *Glob. Environ. Res.* 5: 135-149.

Katoh K S Sakai, Takahashi T. 2009. Factors maintaining species diversity in satoyama, a traditional agricultural landscape of Japan. *Biol. Conserv.* 142: 1930-1936.

Kromp B. 1999. Carabid beetles in sustainable agriculture: a review on pest control efficacy, cultivation impacts and enhancement. *Agric. Ecosyst. Environ.*, 74: 187-228.

- Kwon YJ, SM. 1984. Classification of the subfamily Carabidae from Korea (Coleoptera: Carabidae). *Insecta Koreana*, Series 4, Editorial Committee of Insect Koreana, Seoul, Korea. 149 pp. 107 plates.
- Luff ML, Eyre MD, and Rushton SP. 1989. Classification and ordination of habitats of ground beetles (Coleoptera, Carabidae) in north-east England. *J. Biogeogr.* 16: 121-130.
- Löbl I, Smetana A. 2003. Catalogue of Palaearctic Coleoptera, vol. 1: Archostemata-Myxophaga-Adephaga. Apollo Books, Stenstrup. 819 pp.
- Lövei GL, Sunderland KD. 1996. Ecology and behavior of ground beetles (Coleoptera: Carabidae). *Ann. Rev. Entomol.* 41(1): 231-256.
- Maclean L, Dawe DC, Hardy B, and Hettel GP. 2002. *Rice Almanac*. 3rd ed. International Rice Research Institute, Los Baños.
- Margalef R. 1958. Information theory in ecology. *Gen. Sys.* 3: 36-71.
- McNaughton S J. 1967. Relationship among functional properties of California grassland. *Nature* 216: 168-169.
- Paik CH, Lee GH, Kang JG, Jeon YK, Choi MY, and Seo HY. 2009. Plant flora and insect fauna in the fallow paddy fields of Jeonbuk Province. *Korean J. Appl. Entomol.* 48:285-294. (in Korean with English summary)
- Park JK, Paik JC. 2001. Family Carabidae. *Economic Insects of Korea* 12. *Ins. Koreana Suppl.* 19. 170 pp. (in Korean)
- Park JK. 2004. Subfamily Carabinae in Korea (Coleoptera: Carabidae). *Economic Insects of Korea* 23. *Ins. Koreana Suppl.* 30, 99 pp.
- Pielou EC. 1969. Shannon's formula as a measure of specific diversity, its use and misuse. *Amer. Nat.* 100: 463-465.
- Pielou EC. 1975. *Ecology diversity* Wiley, New York.
- Thomas F, Folgarait P, Lavelle P, and Rossi JP. 2004. Soil macrofaunal communities along an abandoned rice field chronosequence in Northern Argentina. *Appl. Soil. Ecol.* 27: 23-29.
- Uematsu Y, Koga T, Mitsuhashi H, Ushimaru A. 2010. Abandonment and intensified use of agricultural land decrease habitats of rare herbs in semi-natural grasslands. *Agric. Ecosyst. Environ.* 135: 304-309.
- Yamada S, Okubo S, Kitagawa Y, and Takeuchi K. 2007. Restoration of weed communities in abandoned rice paddy fields in the Tama Hills, central Japan. *Agric. Ecosyst. Environ.* 119: 88-102.

Received: 8. Apr. 2013

Revised: 6. May 2013

Accepted: 7. May 2013